



The effect of manual therapy and neuroplasticity education on chronic low back pain: a randomized clinical trial

Adriaan Louw^a, Kevin Farrell^b, Merrill Landers^c, Martin Barclay^b, Elise Goodman^b, Jordan Gillund^b, Sara McCaffrey^b  and Laura Timmerman^b 

^aInternational Spine and Pain Institute, Story City, IA, USA; ^bDepartment of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA; ^cDepartment of Physical Therapy, School of Allied Health Sciences, University of Nevada, Las Vegas, NV, USA

ABSTRACT

Objective: To determine if a neuroplasticity educational explanation for a manual therapy technique will produce a different outcome compared to a traditional mechanical explanation.

Methods: Sixty-two patients with chronic low back pain (CLBP) were recruited for the study. Following consent, demographic data were obtained as well as pain ratings for low back pain (LBP) and leg pain (Numeric Pain Rating Scale), disability (Oswestry Disability Index), fear-avoidance (Fear-Avoidance-Beliefs Questionnaire), forward flexion (fingertips-to-floor), and straight leg raise (SLR) (inclinometer). Patients were then randomly allocated to receive one of two explanations (neuroplasticity or mechanical), a manual therapy technique to their lumbar spine, followed by post-intervention measurements of LBP, leg pain, forward flexion, and SLR.

Results: Sixty-two patients (female 35 [56.5%]), with a mean age of 60.1 years and mean duration of 9.26 years of CLBP participated in the study. There were no statistically significant interactions for LBP ($p = .325$), leg pain ($p = .172$), and trunk flexion ($p = .818$) between the groups, but SLR showed a significant difference in favor of the neuroplasticity explanation ($p = .041$). Additionally, the neuroplasticity group were 7.2 times (95% confidence interval = 1.8–28.6) more likely to improve beyond the MDC on the SLR than participants in the mechanical group.

Discussion: The results of this study show that a neuroplasticity explanation, compared to a traditional biomechanical explanation, resulted in a measureable difference in SLR in patients with CLBP when receiving manual therapy. Future studies need to explore if the increase in SLR correlated to changes in cortical maps of the low back.

KEYWORDS

Pain; brain; plasticity; education; manual therapy; straight leg raise; remapping

Introduction

In the last two decades, the profession of physical therapy (PT), including manual therapy, has seen a significant increase in the interest in pain science.[1,2] Original pain science interest centered on the understanding of pain biology and physiology,[3,4] clinical reasoning incorporating pain mechanisms[4–6] and ultimately, pain neuroscience education.[7–9] By far, the vast majority of this research has focused on chronic low back pain (CLBP).[8–10] In recent years, with the advances in brain scan technology,[11] the focus has shifted toward a greater understanding of the structural and functional changes in the brain of someone suffering from CLBP.[12]

It is now well established that the physical body of a person is represented in the brain by a network of neurons.[11,13–15] This representation refers to the pattern of activity that is evoked when a particular body part is stimulated. The most famous area of the brain associated with representation is the primary somatosensory cortex (S1).[11,13–15] What is interesting is that these neuronal

representations of body parts are dynamically maintained and it has been shown that patients with chronic pain display different S1 representations than people with no pain.[12,16–20] The interesting phenomenon associated with cortical restructuring is the fact that the body maps expand or contract, in essence increasing or decreasing the body map representation in the brain. Furthermore, these changes in shape and size of body maps seem to correlate to increased pain and disability.[12,21] Various factors have been linked to the development of this altered cortical representation of body maps in S1 including neglect and decreased use of the painful body part.[22] This reorganization of body maps occurs quickly. It has been shown that when four fingers are webbed together for 30 min, cortical maps associated with the fingers change.[15] This finding has significant clinical importance as it underscores the importance of strategies such as movement, tactile, and visual stimulation of the central nervous system and brain as a means to help maintain S1 representation.[23,24]

CONTACT Adriaan Louw  ALouw@AOL.com

 The supplemental material for this paper is available online at <https://doi.org/10.1080/10669817.2016.1231860>




Figure 1. Study layout.

In manual therapy, traditional biomechanical models, used to explain to patients a proposed treatment or efficacy of a certain technique or approach, have focused heavily on biomechanical and anatomical models. [25–28] These models would imply that injury, disease, and muscle guarding may lead to altered movement patterns, asymmetrical loading, and resultant pain and dysfunction. [25–28] These biomechanical models have come under scrutiny, partly due to the advances in other fields such as use of diagnostic ultrasound, spinal imaging, and brain scans. [29–31] These advances, with the quest of understanding how a manual technique such as a spinal manipulation or mobilization results in increased range of motion or decreased pain and disability, have led to exciting ‘new’ areas of endeavor. For example, it is now well understood that immediately following a lumbar spine manipulation in someone with low back pain (LBP), there is an immediate neurophysiological effect on the muscles in the affected area. [32,33] Additionally, manual techniques result in endogenous mechanisms in the central nervous system and brain, thus mediating the pain experience. [31,34,35] Findings like these are exciting new frontiers to understand how and why a manual therapy approach helps people in pain. Given our knowledge of rapidly changing functional maps of the human



body in pain, especially CLBP, it could be argued that a manual therapy technique may indeed also be seen as a form of sensory retraining, [23,24] apart from the neuromuscular and endogenous effects previously described. For example, in a recent case series of patients with CLBP, researchers asked patients to identify ‘where they were being touched’ on their low back (for 5 min), and immediately following the sensory discrimination session, the group’s mean pain rating for LBP decreased by 1.91, while forward flexion improved by 4.82 cm, exceeding the minimal detectable change (MDC) of 4.5 cm. [24] The aim of this study was to see if such a neuroplasticity explanation, compared to a traditional biomechanical explanation, would result in superior results in patients with CLBP in regards to pain and movement.

Methodology

Study design

Institutional Review Board approval was obtained for the study. The study was registered as a clinical trial (NCT02757378). This was a randomized clinical trial (RCT) where participants were randomly assigned to either an experimental group (EG) or control group (CG) (Figure 1). All patients acknowledged their understanding and

Table 1. Biomechanical and anatomical explanation of manual therapy.

Neuroplasticity (EG)	Biomechanical (CG)
<p><i>Explanation</i></p> <ul style="list-style-type: none"> • Have a look at this picture – it’s a picture of a brain map of a human body • In your brain there is a map telling you where your body parts are. For example, if we had you close your eyes and touch your nose with your right index finger, you’d have no problem doing it • When life is good and we move during daily tasks, work, and exercise these maps are ‘exercised’ and they stay healthy – sharp and crisp – so we know where the body parts are • When we have pain, move less and do less, the brain areas are not exercised and in essence become blurred • Scientists have now shown us that this happens very fast and the more ‘blurred’ the area is, the more pain we have • We can retrain the brain maps • Today I am going to do some manual treatments to your back as a means to help your brain sharpen its maps 	<p><i>Explanation</i></p> <ul style="list-style-type: none"> • Here is a picture of your low back • There are five bones in your lower back • When life is good and we move, for example, bending forward, each level takes part in the movement and in essence shares the load • When we develop back pain – some levels stiffen up due to swelling and muscle spasms as a means to protect you • I am going to do some treatment on your back with my hands to loosen up your back with the aim to make each level move
<p><i>Picture</i></p>  <p style="text-align: center;">Somatosensory cortex</p>	<p><i>Picture</i></p> 
<p><i>Words during the treatment</i></p> <ul style="list-style-type: none"> • Let the patient know which level you’re on (i.e. L5) and have them verbalize it • When moving to another level, repeat the process 	<p><i>Words during the treatment</i></p> <ul style="list-style-type: none"> • No mention of what is found, but rather a ‘general’ stiffness and manual loosening up each level

willingness to participate by providing signed consent. Participants were informed that the study aimed to determine the efficacy of manual therapy for CLBP.

Randomization

Randomization was performed, using an alternating envelope system. Upon presenting at the clinical site, PTs drew an envelope which randomly assigned the patient to either the EG or CG. The envelopes contained identical information, except that the description of the manual therapy was different for the EG and CG.

Setting

All five PT’s collecting data were orthopedic residents. Each PT treated patients with CLBP, working in separate private practice outpatient clinics in Illinois and Iowa and obtained the necessary permission to participate in the study. Each PT was educated on the aims of the study, measurement tools used in the study, completion of forms, and delivery of the EG and CG words, prior to performing the manual therapy treatment. To ensure uniform application of the manual therapy, each PT, as a resident, reviewed the techniques while in class with a senior manual therapy instructor.

Participants and recruitment

The PTs screened all new patients with CLBP attending PT against the inclusion and exclusion criteria for participation in the study. Inclusion criteria were: adults over the age of 18; presenting at PT with a primary complaint of LBP; LBP being present for 6 months or more; fluent in English; and willing to participate in the study. Exclusion criteria included: medical precautions to the use of manual therapy (metal, skin lesions, etc.); prior spine surgery; and unable to lay prone for the treatment. The sample size was estimated using the Repeated Measures Analysis module in PASS 14 using LBP and a factorial design with one between factor (treatment and control) and one within factor (time). It was based on a Greenhouse-Geisser Corrected F Test for the interaction between the two variables. In order to detect an effect size difference of .45 between the two arms, a total of 48 participants (24 participants in each arm) were needed. The analysis was based on a two-sided test with the significance level at .05 and power at 70%.

Intervention

All patients received the same manual therapy techniques (central posterior–anterior mobilization), position

Table 2. Demographic information.

Characteristic	Results
Age (years; mean)	60.14
Female (%)	35 (56.5%)
White, non-Hispanic	57 (91.9%)
Duration of LBP (years; mean)	9.26
Disability (ODI; mean)	35.67
Fear-avoidance – work subscale (FABQ-W; mean)	17.28
Fear-avoidance – physical activity (FABQ – PA; mean)	16.12
Have received manual therapy before for LBP (n; %)	49 (79%)

(prone), duration (10 min) and grading of the manual technique (grade II), per Maitland.[36] The intervention tested in this RCT was the difference in the educational models used to explain the manual therapy techniques. The descriptions and explanations of the proposed techniques were different between the EG and CG (Table 1) but were equal in duration (5 min).

Outcome measures

Prior to the manual therapy techniques, all study subjects completed a demographics survey capturing age, gender, ethnicity, income, and duration of LBP. Additionally, subjects were asked in regards to any previous exposure to manual therapy and their perceived benefit from previous manual therapy (Likert scale). All patients additionally completed an Oswestry Disability Index (ODI) [37–39] and Fear-Avoidance Beliefs Questionnaire (FABQ) [40] to ascertain their level of disability and fear-avoidance at the time of enrollment into the study. Four measurements were taken prior to, and immediately following manual therapy techniques to determine the efficacy of the experimental intervention:

- *Pain*: LBP and leg pain (if present) was measured using Numeric Pain Rating Scale (NPRS), as has been used in various spinal pain studies.[8,9,41] The MDC for the NPRS for LBP is reported to be 2.0.[42]
- *Lumbar flexion*: Active trunk forward flexion, measured from the longest finger on the dominant hand to the floor.[10,43,44] MDC for active trunk forward flexion has been reported as 4.5 cm.[45]
- *Straight leg raise (SLR)*: We used the SLR as a neurodynamic measurement rather than a test of hamstring length. SLR was measured with an inclinometer placed on the tibial crest 5 cm distal to the

inferior border of the patella on the most affected leg.[10,43,44] SLR for this study kept the ankle in neutral (90°) with no added dorsiflexion or plantar flexion, per previous studies.[10,43,44] MDC for SLR has been reported as a 5.7° difference.[45]

Patients completed the NPRS, underwent lumbar flexion and SLR testing immediately prior to receiving the manual therapy technique and immediately after manual therapy technique. Pre- and post-treatment measurements were performed by the therapists who provided the manual interventions. Following capture of the outcome measures and manual therapy, the PTs continued their treatment per their plan of care.

Statistical analysis

In order to ascertain any differences between the treatment and control, 2 (group: treatment and control) × 2 (time: pre and post) mixed factorial ANOVAs on four different outcome measures (LBP, leg pain, trunk flexion, SLR) were conducted. If an interaction between group and time was observed then simple main effects were conducted using a Bonferroni correction of .025 for two simple main effects tests for the within analysis (pre and post) of each group. If no interaction was observed, then main effects were analyzed. Chi-square analysis was used to compare the proportions of individuals in each treatment who exceeded the MDC for each dependent measure.

Results

Participants

Sixty-two patients (female 35 [56.5%]), with a mean age of 60.14 years and mean duration of 9.26 years of LBP, participated in the study (Table 2). A majority of the patients ($n = 49$ [79%]) had received manual therapy previously for their LBP. In general, patients had a positive experience of manual therapy for LBP, with an average score of 5.97/10 agreeing manual therapy had helped their LBP before (0 – do not agree; 10 – strongly agree).

Outcome measures

There were no statistically significant interactions for LBP ($p = .325$), leg pain ($p = .172$), and trunk flexion ($p = .818$)

Table 3. Means with standard deviations, including main effects of group and time, for each of the outcome variables.

Variable	Group	Pre – means and SDs	Post – means and SDs	Interaction effect	Time main effect	Group main effect
LBP	Treatment	3.8 ± 2.1	3.0 ± 2.4	$p = .325$	$p = .004^*$	$p = .180$
	Control	4.3 ± 2.4	4.0 ± 2.5			
Leg pain	Treatment	2.8 ± 2.4	2.1 ± 2.5	$p = .172$	$p = .094$	$p = .892$
	Control	2.6 ± 2.6	2.5 ± 2.2			
Trunk flexion	Treatment	25.9 ± 19.1	23.1 ± 19.8	$p = .818$	$p = .004^*$	$p = .853$
	Control	25.3 ± 15.4	22.1 ± 16.2			
SLR	Treatment	51.2 ± 18.1	56.2 ± 19.0	$p = .041^*$	$p < .001^*$	$p = .662$
	Control	55.2 ± 20.7	56.6 ± 21.0			

*Statistically significant at $p < .05$.




Figure 2. Means and standard errors of the mean for the straight leg raise at the pre- and post-measurements for both the experimental and control groups.

Table 4. 2 × 2 contingency table for the proportion of participants who improved beyond the MDC on the SLR.

		Yes		No	
Group	Treatment	Count	15	18	
		% within group	45.5	54.5	
	Control	Count	3	26	
		% within group	10.3	89.7	
Total		Count	18	44	
		% within group	29.0	71.0	

(Table 3). However, there was a significant interaction for SLR ($p = .041$) (Table 3, Figure 2). Simple main effects analyses indicate that the SLR improved for the EG ($p = .001$) but not for the CG ($p = .123$) (Figure 2). Descriptive data and main effect p values for each outcome variable are available in Table 3.

The proportion of participants across the two groups who improved beyond the MDC was compared using chi-square analyses (Fisher's exact test). There were no statistically significant differences for LBP ($p = .055$), leg pain ($p = .676$), and trunk flexion ($p = 1.000$); however, there was for SLR ($p = .004$). Participants in the treatment group were 7.2 times (95% confidence interval for odds ratio = 1.8–28.6) more likely to improve beyond the MDC on the SLR than participants in the CG (Table 4).

Discussion

The results of this study show that a neuroplasticity explanation, compared to a traditional biomechanical explanation, resulted in a measurable difference in the likelihood of improved SLR response in patients with CLBP when receiving manual therapy.

In the current study, with similar manual techniques applied to patients with CLBP, yet receiving different educational models for the manual techniques resulted in a difference in SLR, exceeding the MDC.[45] These results concur with various other pain science studies whereby pain neuroscience education resulted in immediate changes in neurodynamic tests such as SLR and the upper limb neurodynamic tests.[2,44,46–48] Coppieters et al. showed that altering the explanation of the SLR test by itself from describing the test as a test of 'muscle' vs. 'nerve' also significantly alters SLR, showcasing once

again the power of words altering SLR.[49] In contrast, neither explanation resulted in a significant shift in LBP, leg pain or forward flexion. The results in regards to pain are expected. Given the complexity of pain, especially chronic pain,[1,50] it is not expected an education-alone session should result in immediate meaningful changes. This assumption is based on various pain neuroscience education studies showing no immediate effect on pain alone.[51–53] There is even evidence for an educational phenomenon referred to as 'explain pain pain,' whereby education about pain neuroscience results in a slight increase in pain after education, followed by a decrease over time.[47,54]

In regards to forward flexion, previous pain neuroscience education studies showed immediate changes in forward flexion[44,47,54] but not the current study. This study, however, was not a pain neuroscience study, but rather a comparison of different explanations for a manual therapy technique. In pain neuroscience education, a more elaborate discussion of pain science is undertaken.[55,56] The fact that SLR changed in the short amount of time and not flexion is intriguing. One potential explanation may be that SLR changes may be associated with the fact that neurodynamics is likely more affected by processes which may rapidly respond to education such as blood supply, muscle tone, etc.[57–60] In contrast, forward flexion, a common physical examination,[61] and often associated with LBP may need more extensive education, including addressing fear of movement.[61] In the current study, the mean FABQ-PA was 16.12, and an FABQ-PA > 14 is associated with a higher likelihood of not returning to work,[62,63] which would imply a 'simple' educational explanation for a manual technique would not be effective to alter flexion, but likely warrant a more comprehensive educational session.

The current study did not aim to specifically examine if cortical changes occurred in line with the education and manual techniques. Future studies will be needed to examine if these techniques and resultant increased SLR were in fact associated with cortical remapping. The results, however, do add to the body of growing evidence that manual therapy can be seen as a form of sensory discrimination, integration, and cortical remapping. For example, recent pain neuroscience education studies using single-case functional magnetic resonance imaging (fMRI) [41,47] have shown immediate cortical changes in the brain of patients with LBP, with one showing an immediate change in the motor cortex, as well as the patient's SLR improving 7° following the education.[47] Other studies aiming to examine if tactile information via manual touch has a therapeutic effect in LBP, have shown to result in SLR changes,[23] immediate hypoalgesic effect,[23] and improved forward flexion exceeding the MDC.[24] Additionally, it has been shown there are immediate cortical changes in the brain following thoracic manipulation [64] as well as motor control exercises.[65,66]

Although a clear picture is yet to emerge in regards to manual therapy, cortical remapping and its influence on pain, function and movement, it does underscore the need for pain science to be hands-on vs. hands-off.[67] The interest in pain neuroscience education has likely driven the pendulum away from traditional manual approaches, which may in fact be detrimental.[68] A new systematic review on the efficacy of pain neuroscience education has highlighted that pain-education sessions when combined with movement result in superior results when compared to education alone.[56] The current study underscores the need for combining education with manual therapy, ultimately strengthening the therapeutic alliance, which is defined as the working rapport or positive social connection between the patient and the therapist.[69–71] By virtue of its definition, therapeutic alliance is a complex blend of therapist technical skill, verbal and non-verbal communication, sense of warmth, trust, and collaboration.[69] One factor, heavily associated with therapeutic alliance, is the words clinicians chose in explaining a test or treatment. It is well established that word-choice by health care providers can have a positive or negative influence on their patients, which concurs with the current study.

This study contains various limitations. First, as described, the increased SLR cannot be directly attributed to cortical reorganization. The treatment techniques (Grade II) were arbitrarily chosen to not agitate the patient's condition and not matched to presenting signs and symptoms, let alone specificity of the technique for the condition (a central posterior–anterior mobilization vs. other techniques). Even with an attempt to standardize the delivery of the education and manual therapy, there would have been differences in experience with the therapists delivering the treatment. A significant limitation is the fact that tests and treatments were performed with the attending therapists, thus the testing therapists were not blinded to the treatment. No attempt was made to differentiate if the education before the treatment or education during the treatment was responsible for the proposed change, or the combination of the two educational sessions, and future research should further explore this possibility. It is also important to point out that the MDC may not be clinically important or relevant; however, no minimal clinically important difference value has been reported for the SLR.[72,73]

Conclusion

The results of this study indicate that a neuroplasticity explanation, compared to a traditional biomechanical explanation, resulted in a measureable difference in SLR in patients with CLBP. The results also indicate no differences between the groups in regards to LBP, leg pain, and forward flexion. Future studies should investigate

long-term effects of such education and may be also from a combination of therapies including neuroscience education.

Acknowledgements

The author Adriaan Louw, PT, PhD, acknowledges receiving payment for various speaking events (seminars, conferences, etc.) as well as royalties from books, from professionals and patients.

Disclosure statement

None of these conflicts influence the subject matter or materials discussed in this manuscript.

Notes on contributors

Adriaan Louw, PT, PhD is the owner and CEO of International Spine and Pain Institute, Story City, IA, USA. He is an adjunct faculty in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA. He is an adjunct faculty: Department of Physical Therapy, School of Allied Health Sciences, University of Nevada, Las Vegas, NV, USA. He is the program director of Evidence in Motion Pain Certification and Pain Fellowship.

Kevin Farrell, PT, PhD, OCS, FAAOMPT, is a professor and chair of the Orthopaedic Clinical Residency in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

Merrill Landers, PT, PhD, DPT, OCS, is the chair and professor in the Department of Physical Therapy, School of Allied Health Sciences, University of Nevada, Las Vegas, NV, USA. He is also Cyrus Chung Ying Tang Foundation Research Professor and fellow of the APTA Education Leadership Institute.

Martin Barclay, PT, DPT, OCS is a Resident in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

Elise Goodman, PT, DPT, OCS is a resident in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

Jordan Gillund, PT, DPT, OCS is a resident in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

Sara McCaffrey, PT, DPT, OCS is a resident in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

Laura Timmerman, PT, DPT, OCS is a resident in the Department of Physical Therapy Education, Residency Program, St. Ambrose University, Davenport, IA, USA.

ORCID

Sara McCaffrey  <http://orcid.org/0000-0002-8956-814X>

Laura Timmerman  <http://orcid.org/0000-0001-8162-500X>

References

- [1] Moseley GL. Reconceptualising pain according to modern pain science. *Phys. Ther. Rev.* 2007;12:169–178.
- [2] Nijs J, Roussel N, Paul van Wilgen C, et al. Thinking beyond muscles and joints: therapists' and patients' attitudes and beliefs regarding chronic musculoskeletal pain are key to applying effective treatment. *Man. Ther.* 2013;18:96–102.
- [3] Gifford LS. Pain, the tissues and the nervous system: a conceptual model. *Physiotherapy.* 1998;84:27–36.
- [4] Gifford L, Butler D. The integration of pain sciences into clinical practice. *J. Hand Ther.* 1997;10:86–95.
- [5] Jones MA. Clinical reasoning: the foundation of clinical practice. Part 1. *Aust. J. Physiother.* 1997;43:167–170.
- [6] Jones M. Clinical reasoning and pain. *Man. Ther.* 1995;1:17–24.
- [7] Gifford L, Muncey H. Explaining pain to patients. Paper presented at: International Association on the Study of Pain 1999; Vienna, Austria.
- [8] Moseley L. Combined physiotherapy and education is efficacious for chronic low back pain. *Aust. J. Physiother.* 2002;48:297–302.
- [9] Moseley GL. Joining forces – combining cognition-targeted motor control training with group or individual pain physiology education: a successful treatment for chronic low back pain. *J. Man. Manipulative Ther.* 2003;11:88–94.
- [10] Moseley GL, Hodges PW, Nicholas MK. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin. J. Pain.* 2004;20:324–330.
- [11] Flor H. The functional organization of the brain in chronic pain. *Prog. Brain Res.* 2000;129:313–322.
- [12] Flor H, Braun C, Elbert T, et al. Extensive reorganisation of primary somatosensory cortex in chronic back pain patients. *Neurosci. Lett.* 1997;224:5–8.
- [13] Wand BM, Parkitny L, O'Connell NE, et al. Cortical changes in chronic low back pain: current state of the art and implications for clinical practice. *Man. Ther.* 2011;16:15–20.
- [14] Penfield W, Boldrey E. Somatic, motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain.* 1937;60:389–443.
- [15] Stavrinou ML, Della Penna S, Pizzella V, et al. Temporal dynamics of plastic changes in human primary somatosensory cortex after finger webbing. *Cereb. Cortex.* 2007;17:2134–2142.
- [16] Maihofner C, Handwerker HO, Neundorfer B, et al. Patterns of cortical reorganization in complex regional pain syndrome. *Neurology.* 2003;61:1707–1715.
- [17] Moseley GL. I can't find it! Distorted body image and tactile dysfunction in patients with chronic back pain. *Pain.* 2008;140:239–243.
- [18] Lotze M, Moseley GL. Role of distorted body image in pain. *Curr. Rheumatol. Rep.* 2007;9:488–496.
- [19] Moseley GL. Distorted body image in complex regional pain syndrome. *Neurology.* 2005;65:773.
- [20] Flor H, Elbert T, Muhnckel W, et al. Cortical reorganisation and phantom phenomena in congenital and traumatic upper-extremity amputees. *Exp. Brain Res.* 1998;119:205–212.
- [21] Lloyd D, Findlay G, Roberts N, et al. Differences in low back pain behavior are reflected in the cerebral response to tactile stimulation of the lower back. *Spine.* 2008;33:1372–1377.
- [22] Marinus J, Moseley GL, Birklein F, et al. Clinical features and pathophysiology of complex regional pain syndrome. *Lancet Neurol.* 2011;10:637–648.
- [23] Louw A, Schmidt SG, Louw C, et al. Moving without moving: immediate management following lumbar spine surgery using a graded motor imagery approach: a case report. *Physiother. Theory Pract.* 2015;31:509–517.
- [24] Louw A, Farrell K, Wettach L, et al. Immediate effects of sensory discrimination for chronic low back pain: a case series. *N. Z. J. Physiother.* 2015;43:58–63.
- [25] Twomey LT. A rationale for the treatment of back pain and joint pain by manual therapy. *Phys. Ther.* 1992;72:885–892.
- [26] Edmondston SJ, Singer KP. Thoracic spine: anatomical and biomechanical considerations for manual therapy. *Man. Ther.* 1997;2:132–143.
- [27] Edmondston SJ, Allison GT, Gregg CD, et al. Effect of position on the posteroanterior stiffness of the lumbar spine. *Man. Ther.* 1998;3:21–26.
- [28] Barrett CJ, Singer KP, Day R. Assessment of combined movements of the lumbar spine in asymptomatic and low back pain subjects using a three-dimensional electromagnetic tracking system. *Man. Ther.* 1999;4:94–99.
- [29] Flynn TW. There's more than one way to manipulate a spine. *J. Orthop. Sports Phys. Ther.* 2006;36:198–199.
- [30] Flynn TW. Manual physical therapy: moving beyond the theoretical. *J. Orthop. Sports Phys. Ther.* 2004;34:659–661.
- [31] Bialosky JE, Bishop MD, Price DD, et al. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man. Ther.* 2009;14:531–538.
- [32] Raney NH, Teyhen DS, Childs JD. Observed changes in lateral abdominal muscle thickness after spinal manipulation: a case series using rehabilitative ultrasound imaging. *J. Orthop. Sports Phys. Ther.* 2007;37:472–479.
- [33] Puentedura EJ, Landers MR, Hurt K, et al. Immediate effects of lumbar spine manipulation on the resting and contraction thickness of transversus abdominis in asymptomatic individuals. *J. Orthop. Sports Phys. Ther.* 2011;41:13–21.
- [34] Bialosky JE, Bishop MD, George SZ, et al. Placebo response to manual therapy: something out of nothing? *J. Man. Manipulative Ther.* 2011;19:11–19.
- [35] Bialosky JE, Bishop MD, Robinson ME, et al. Spinal manipulative therapy has an immediate effect on thermal pain sensitivity in people with low back pain: a randomized controlled trial. *Phys. Ther.* 2009;89:1292–1303.
- [36] Maitland GD. *Vertebral manipulation.* 6th ed. London: Butterworths; 1986.
- [37] Deyo RA, Battie M, Beurskens AJ, et al. Outcome measures for low back pain research: a proposal for standardized use. *Spine.* 1998;23:2003–2013.
- [38] Fritz JM, Irrgang JJ. A comparison of a modified Oswestry low back pain disability questionnaire and the Quebec back pain disability scale. *Phys. Ther.* 2001;81:776–788.
- [39] Hakkinen A, Kautiainen H, Jarvenpaa S, et al. Changes in the total Oswestry index and its ten items in females and males pre- and post-surgery for lumbar disc herniation: a 1-year follow-up. *Eur. Spine J.* 2007;16:347–352.
- [40] George SZ, Fritz JM, Childs JD. Investigation of elevated fear-avoidance beliefs for patients with low back pain: a secondary analysis involving patients enrolled in physical therapy clinical trials. *J. Orthop. Sports Phys. Ther.* 2008;38:50–58.
- [41] Moseley GL. Widespread brain activity during an abdominal task markedly reduced after pain physiology education: fMRI evaluation of a single patient with chronic low back pain. *Aust. J. Physiother.* 2005;51:49–52.
- [42] Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine.* 2005;30:1331–1334.

- [43] Zimney K, Louw A, Puentedura EJ. Use of therapeutic neuroscience education to address psychosocial factors associated with acute low back pain: a case report. *Physiother. Theory Pract.* 2014;30:202–209.
- [44] Moseley GL. Evidence for a direct relationship between cognitive and physical change during an education intervention in people with chronic low back pain. *Eur. J. Pain.* 2004;8:39–45.
- [45] Ekedahl H, Jonsson B, Frobell RB. Fingertip-to-floor test and straight leg raising test: validity, responsiveness, and predictive value in patients with acute/subacute low back pain. *Arch. Phys. Med. Rehabil.* 2012;93:2210–2215.
- [46] Louw A, Puentedura EL, Mintken P. Use of an abbreviated neuroscience education approach in the treatment of chronic low back pain: a case report. *Physiother. Theory Pract.* 2012;28:50–62.
- [47] Louw A, Puentedura EJ, Diener I, et al. Preoperative therapeutic neuroscience education for lumbar radiculopathy: a single-case fMRI report. *Physiother. Theory Pract.* 2015;31:496–508.
- [48] Meeus M, Nijs J, Van Oosterwijck J, et al. Pain physiology education improves pain beliefs in patients with chronic fatigue syndrome compared with pacing and self-management education: a double-blind randomized controlled trial. *Arch. Phys. Med. Rehabil.* 2010;91:1153–1159.
- [49] Coppieters MW, Ryan L, Chan KP, et al. Do patients' beliefs based on widespread medical information hinder accurate diagnosis? Paper Presented at: 11th World Congress on Pain2005; Sydney.
- [50] Melzack R. Pain and the neuromatrix in the brain. *J. Dent. Educ.* 2001;65:1378–1382.
- [51] Louw A, Diener I, Landers MR, et al. Preoperative pain neuroscience education for lumbar radiculopathy: a multicenter randomized controlled trial with 1-year follow-up. *Spine.* 2014;39:1449–1457.
- [52] van Ittersum MW, van Wilgen CP, van der Schans CP, et al. Written pain neuroscience education in fibromyalgia: a multicenter randomized controlled trial. *Pain Pract.* 2014;14:689–700.
- [53] Ryan CG, Gray HG, Newton M, et al. Pain biology education and exercise classes compared to pain biology education alone for individuals with chronic low back pain: a pilot randomised controlled trial. *Man. Ther.* 2010;15:382–387.
- [54] Louw A, Diener I, Puentedura EJ. The short term effects of preoperative neuroscience education for lumbar radiculopathy: a case series. *Int. J. Spine Surg.* 2015;9:11.
- [55] Louw A, Diener I, Butler DS, et al. The effect of neuroscience education on pain, disability, anxiety, and stress in chronic musculoskeletal pain. *Arch. Phys. Med. Rehabil.* 2011;92:2041–2056.
- [56] Louw A, Zimney K, Puentedura E, et al. The efficacy of pain neuroscience education on musculoskeletal pain: A systematic review of the literature. *Physiother. Theory Pract.* 2016;Jun 28:1–24.
- [57] von Piekartz HJ, Schouten S, Aufdemkampe G. Neurodynamic responses in children with migraine or cervicogenic headache versus a control group: a comparative study. *Man. Ther.* 2007;12:153–160.
- [58] Nee RJ, Butler D. Management of peripheral neuropathic pain: integrating neurobiology, neurodynamics and clinical evidence. *Phys. Ther. Sport.* 2006;7:36–49.
- [59] Coppieters MW, Butler DS. Do 'sliders' slide and 'tensioners' tension? An analysis of neurodynamic techniques and considerations regarding their application. *Man. Ther.* 2008;13:213–221.
- [60] Nee RJ, Vicenzino B, Jull GA, et al. Neural tissue management provides immediate clinically relevant benefits without harmful effects for patients with nerve-related neck and arm pain: a randomised trial. *J. Physiother.* 2012;58:23–31.
- [61] George SZ, Fritz JM, McNeil DW. Fear-avoidance beliefs as measured by the fear-avoidance beliefs questionnaire: change in fear-avoidance beliefs questionnaire is predictive of change in self-report of disability and pain intensity for patients with acute low back pain. *Clin. J. Pain.* 2006;22:197–203.
- [62] Fritz JM, George SZ. Identifying psychosocial variables in patients with acute work-related low back pain: the importance of fear-avoidance beliefs. *Phys. Ther.* 2002;82:973–983.
- [63] Burton AK, Waddell G, Tillotson KM, et al. Information and advice to patients with back pain can have a positive effect: a randomized controlled trial of a novel educational booklet in primary care. *Spine.* 1999;24:2484–2491.
- [64] Sparks C, Cleland JA, Elliott JM, et al. Using functional magnetic resonance imaging to determine if cerebral hemodynamic responses to pain change following thoracic spine thrust manipulation in healthy individuals. *J. Orthop. Sports Phys. Ther.* 2013;43:340–348.
- [65] Tsao H, Galea MP, Hodges PW. Reorganization of the motor cortex is associated with postural control deficits in recurrent low back pain. *Brain.* 2008;131:2161–2171.
- [66] Tsao H, Danneels LA, Hodges PW. ISSLS prize winner: smudging the motor brain in young adults with recurrent low back pain. *Spine.* 2011;36:1721–1727.
- [67] Lluch Girbes E, Meeus M, Baert I, et al. Balancing, "hands-on" with "hands-off" physical therapy interventions for the treatment of central sensitization pain in osteoarthritis. *Man. Ther.* 2015;20:349–352.
- [68] Louw A, Puentedura E, Zimney K, et al. Know pain; know gain? A perspective on pain neuroscience education in physical therapy. *J. Orthop. Sports Phys. Ther.* 2016;46:131–134.
- [69] Crepeau EB, Garren KR. I looked to her as a guide: the therapeutic relationship in hand therapy. *Disabil. Rehabil.* 2011;33:872–881.
- [70] Fuentes J, Armijo-Olivo S, Funabashi M, et al. Enhanced therapeutic alliance modulates pain intensity and muscle pain sensitivity in patients with chronic low back pain: an experimental controlled study. *Phys. Ther.* 2014;94:477–489.
- [71] Joyce AS, Ogradniczuk JS, Piper WE, et al. The alliance as mediator of expectancy effects in short-term individual therapy. *J. Consult. Clin. Psychol.* 2003;71:672–679.
- [72] Dixon JK, Keating JL. Variability in straight leg raise measurements. *Physiotherapy.* 2000;86:361–370.
- [73] Boyd BS. Measurement properties of a hand-held inclinometer during straight leg raise neurodynamic testing. *Physiotherapy.* 2012;98:174–179.